

New records of bees of the genus *Sphecodes* Latreille in the Palaearctic part of China (Hymenoptera, Halictidae)

Yulia V. Astafurova¹, Maxim Yu. Proshchalykin², Ze-qing Niu³, Chao-dong Zhu^{3,4}

1 Zoological Institute, Russian Academy of Sciences, Universitetskaya Nab., 1, Saint Petersburg 199034, Russia
2 Federal Scientific Centre for East Asian Terrestrial Biodiversity, Far Eastern Branch of Russian Academy of Sciences, Vladivostok 690022, Russia **3** Key Laboratory of Zoological Systematics and Evolution, Institute of Zoology, Chinese Academy of Sciences, 1 Beichen West Road, Chaoyang District, Beijing, 100101, PR China
4 College of Life Sciences, University of Chinese Academy of Sciences, No.19(A) Yuquan Road, Shijingshan District, Beijing, 100049, PR China

Corresponding author: Maxim Yu. Proshchalykin (proshchalikin@biosoil.ru)

Academic editor: M. S Engel | Received 28 June 2018 | Accepted 15 August 2018 | Published 23 October 2018

<http://zoobank.org/B6B988C5-521B-4854-B9B0-1F9713779323>

Citation: Astafurova YV, Proshchalykin MY, Niu Z-q, Zhu C-d (2018) New records of bees of the genus *Sphecodes* Latreille in the Palaearctic part of China (Hymenoptera, Halictidae). ZooKeys 792: 15–44. <https://doi.org/10.3897/zookeys.792.28042>

Abstract

The available information about the cleptoparasitic bees of the genus *Sphecodes* in the Palaearctic part of China is summarized. Twenty-four species are currently known from this area including 16 newly recorded. Based on type specimens, new synonymies have been proposed for *Sphecodes cristatus* Hagens, 1882 = *S. alfkeni* Meyer, 1922, **syn. n.**; *S. longulus* Hagens, 1882 = *S. subfasciatus* Blüthgen, 1934, **syn. n.**; *S. nippon* Meyer, 1922 = *S. kansuensis* Blüthgen, 1934, **syn. n.**; *Sphecodes pieli* Cockerell, 1931 = *S. orientalis* Astafurova & Proshchalykin, 2014, **syn. n.** Lectotypes are designated for *Sphecodes alfkeni* Meyer, 1922 and *S. pellucidus niveipennis* Meyer, 1925. Illustrated keys to males and females of all species known from Palaearctic China and an updated checklist of the 33 Chinese species of *Sphecodes* are provided.

Keywords

Anthophila, Apiformes, cleptoparasites, fauna, new synonymy, taxonomy

Introduction

The present paper is part of a series of works dealing with the bees of the cleptoparastic genus *Sphecodes* Latreille, 1804 from the Palaearctic region (Astafurova and Proshchalykin 2014, 2015a, b, 2016a, b, 2017a, b, Astafurova et al. 2014, 2015, 2018). Consequently, we focus on species in northern China and do not deal with the southern, Oriental species of China.

The question of where the zoogeographical boundary exists between the Oriental and the Palaearctic regions in China has been discussed by many researchers working on various groups of animals (Emeljanov 1974, Hoffman 2001, Fellowes 2006, Chen et al. 2008, Heiser and Schmitt 2013, He et al. 2017). In this paper, the views of Pesenko (2007) and Astafurova (2013) are followed for halictid bees, which posit that the approximate border between Palaearctic and Oriental Regions in China lies between 30°–35° northern latitude.

Sphecodes formosanus Cockerell was the first species of the genus described from China (Taiwan) (Cockerell 1911). Since then, a total of ten species and three subspecies have been described (Meyer, four species and two subspecies; Cockerell, four species; Blüthgen, two species and one subspecies), seven of which are still valid (see section on taxonomy for details). Sixteen *Sphecodes* species have been recorded from China so far (Meyer 1920, 1922, 1925, Blüthgen 1924, 1927, 1934, Cockerell 1911, 1922, 1931, Strand and Yasumatsu 1938, Ascher and Pickering 2018). Among them, only seven species were known from the Palaearctic.

Based on a comprehensive study of specimens in various collections, we catalogue 24 species of the genus *Sphecodes*, with 16 species recorded from China for the first time. New synonymies are proposed for four specific names: *Sphecodes cristatus* Hagens, 1882 = *S. alfkeni* Meyer, 1922, syn. n.; *S. longulus* Hagens, 1882 = *S. subfasciatus* Blüthgen, 1934, syn. n.; *S. nippon* Meyer, 1922 = *S. kansuensis* Blüthgen, 1934, syn. n.; *Sphecodes pieli* Cockerell, 1931 = *S. orientalis* Astafurova & Proshchalykin, 2014, syn. n. Illustrated keys to the species known from the Palaearctic part of China are presented to facilitate further studies.

Materials and methods

The results presented in this paper are based on 453 specimens collected in the Palaearctic part of China and currently housed in the Institute of Zoology, Chinese Academy of Sciences (Beijing, China, IZCAS); the Zoological Institute, Russian Academy of Sciences (St. Petersburg, Russia, ZISP); and the private collection of Maximilian Schwarz (Anselden, Austria, PCMS). The following acronyms are used for the collections where type specimens are deposited:

MNHB Museum für Naturkunde der Humboldt Universität zu Berlin, Germany.

NHRS Naturhistoriska riksmuseet, Stockholm, Sweden.

USNM National Museum of Natural History, Smithsonian Institution, Washington, DC, USA.

KUF Kyushu University, Fukuoka, Japan.

The taxonomy and distribution of species generally follow that of Warncke (1992), Bogusch and Straka (2012), and Astafurova and Proshchalykin (2017b). A detailed current synonymy of the species has been given by Astafurova and Proshchalykin (2017b). Morphological terminology follows that of Michener (2007) and Engel (2001). The ventral surface of some flagellomeres bear a distinctive patch of sensilla trichodea A (sensu Årgent and Svensson 1982), which we refer to as “tyloids;” they are easily observable under light microscopy. The abbreviations F, T, and S are used for flagellomere, metasomal tergum, and metasomal sternum, respectively. The density of integumental punctures is described using the following formula: puncture diameter (in μm) / ratio of distance between punctures to average puncture diameter, e.g., 15–20 μm / 0.5–1.5. Integumental sculpturing, aside from distinctive surface punctuation, is described as follows: reticulate: superficially net-like or made up of a network of raised lines; rugose: irregular, non-parallel, wrinkled raised lines (rugae); tessellate: regular network of shallow grooves with flat interspaces.

Specimens were studied with a Leica M205A stereomicroscope and photographs were taken with a combination of a stereomicroscope (Olympus SZX10) and a digital camera (Canon EOS70D). Final images represent a composite of several photographs taken at different focal planes and combined using the program Helicon Focus 6. All images were post-processed for contrast and brightness using Adobe® Photoshop®.

The species are presented alphabetically and those that could not be inspected in this paper are quoted from published sources. We use the following abbreviations for collectors: **JH** – Jiri Halada, **PK** – Petr Kozlov; **VR** – Vsevolod Roborovsky. New distributional records are noted with an asterisk (*).

Unfortunately, we have not examined the type of *S. manchurianus*, because it was not found in Kyushu University (Japan). We also have not found specimens corresponding to the original description of Strand and Yasumatsu (1938) in our material.

Taxonomy

Key to the *Sphecodes* species of the Palearctic part of China

Additional species are included in this key because they are widespread in the Palearctic and may also occur in China. These include *S. maruyamanus* Tsuneki, 1983, *S. murotai* Tsuneki, 1983, *S. tanoi* Tsuneki, 1983 (known from Japan and Russian Far East), *S. miniatus* Hagens, 1882, *S. puncticeps* Thomson, 1870, and *S. reticulatus* Thomson, 1870. *Sphecodes manchurianus* Strand & Yasumatsu, 1938, known only from the holotype, is not included.

Males

- 1 Costal margin of hind wing with 7–14 hamuli. Base of gonocoxite dorsally without impression. Usually large species: total body length 5.0–12.0 mm**2**
- Costal margin of hind wing with 5–6 hamuli. Base of gonocoxite dorsally with or without impression. Large or small species: total body length 3.5–11.0 mm**11**
- 2 Head rounded, about as long as wide. Hind wing with basal (*M*) vein strongly curved. T1 finely and sparsely (sometimes indistinctly) punctate. Gonostylus dorsally with small rectangular process directed to penis valve (Fig. 15). Body length 7.0–10.0 mm ***S. monilicornis* (Kirby)**
- Head transverse, wider than long. Hind wing with basal (*M*) vein weakly curved. T1 distinctly coarsely and densely punctate. Gonostylus another shape **3**
- 3 Mesoscutum densely punctate, with confluent punctures (areolate)**4**
- Mesoscutum sparsely punctate, medially with punctures separated by at least a puncture diameter **5**
- 4 Head more transverse, 1.2 times as wide as long. Vertex long, distance from top of head to upper margin of lateral ocellus about 2.5–3.0 lateral ocellar diameters as seen in dorsal view. Tyloids on flagellomeres (at least from F4 onward) semicircular across basal 1/5–1/3 and linear across remaining flagellomeres as seen in lateral view. Mesoscutellum sparsely punctate, medially with punctures separated by more than a puncture diameter and often with impunctate areas. T1 completely red. Gonostylus larger, not narrowed apically (Fig. 1). Body length 9.0–12.0 mm ***S. albilabris* (Fabricius)**
- Head less transverse, 1.1 times as wide as long. Vertex shorter, distance from top of head to upper margin of lateral ocellus about two lateral ocellar diameters as seen in dorsal view. Tyloids on flagellomeres semicircular across basal 1/6–1/4, linear portion along remaining flagellomeres not developed. Mesoscutellum densely punctate, with confluent punctures. T1 black or brownish at least on basal 1/3 Gonostylus smaller, distinctly narrowed apically (Fig. 23). Body length 7.0–12.0 mm ***S. scabricollis* Wesmael**
- 5 Vertex with a longitudinal carina. Gonostylus smaller, not overlapped apically, as in Figs 4, 21 **6**
- Vertex without a longitudinal carina. Gonostylus larger, another shape, overlapped apically **8**
- 6 Tyloids on flagellomeres (at least from F4 onward) are semicircular across basal 1/3–1/2. T1 with marginal zone very finely and indistinctly punctate. Body length 7.0–10.0 mm ***S. cristatus* Hagens**
- Tyloids on flagellomeres weakly developed, very narrow, semicircular across basal 1/7–1/5 of flagellomere. T1 with marginal zone coarsely and distinctly punctate **7**

- 7 Head more transverse, 1.2 times as wide as long. Mesoscutum more coarsely punctate (30–75 μm). T2 with marginal zone impunctate. Larger: total body length 7.0–11.0 mm.....***S. olivieri* Lepeletier de Saint-Fargeau**
- Head less transverse, 1.10–1.15 times as wide as long. Mesoscutum more finely punctate (25–40 μm). T2 with marginal zone distinctly punctate. Smaller: total body length 5.0–7.0 mm.....***S. pectoralis* Morawitz**
- 8 Vertex long, distance from top of head to upper margin of lateral ocellus about three lateral ocellar diameters as seen in dorsal view. Tyloids on flagellomeres cover at least 1/3 part of flagellomere Gonostylus with long apical process (Fig. 9).....**9**
- Vertex shorter, distance from top of head to upper margin of lateral ocellus about two lateral ocellar diameters as seen in dorsal view. Tyloids on flagellomeres not covering more than 1/4 part of flagellomere. Gonostylus another shape at tip, as in Fig. 2**10**
- 9 Tyloids on flagellomeres well developed, covering large part of flagellomere (as seen in lateral view, Fig. 57). Body length 7.0–14.0 mm***S. gibbus* (Linnaeus)**
- Tyloids on flagellomeres weakly developed, covering about 1/3 of flagellomere (as seen in lateral view, Fig. 58). Body length 7.0–11.0 mm***S. nippon* Meyer**
- 10 T4 with marginal zone finely tessellate, without punctures (Fig. 46). Body length 7.0–10.0 mm..... ***S. reticulatus* Thomson**
- T4 with marginal zone distinctly punctate, smooth between punctures (rarely indistinctly tessellate) (Fig. 45). Body length 7.0–12.0 mm.....***S. alternatus* Smith**
- 11 Base of gonocoxite dorsally without impression**12**
- Base of gonocoxite dorsally with impression**19**
- 12 T1 densely punctate. Gonostylus elongate (Fig. 18). Body length 5.0–5.5 mm..***S. nurekensis* Warncke**
- T1 impunctate or with a few fine punctures. Gonostylus another shape**13**
- 13 Vertex coarsely and densely punctate, ocello-ocular area with confluent punctures, separated by at most a half puncture diameter**14**
- Vertex finely and sparser punctate, ocello-ocular area with punctures, separated by at least a puncture diameter.....**17**
- 14 Vertex with longitudinal carina (in *S. kozlovi* usually weakly developed) (Figs 38, 39)**15**
- Vertex without longitudinal carina.....**16**
- 15 Vertex with well visible longitudinal carina. Felt-like areas on last flagellomeres cover at least 1/2 underside of flagellomere, F2 as long as wide (Fig. 53). T1 impunctate or with a few fine punctures. Membranous portion of gonostylus smaller, as in Fig. 19. Body length 6.0–9.0 mm..... ***S. pieli* Cockerell**
- Vertex with weakly visible longitudinal carina. Felt-like areas on last flagellomeres cover about 1/3 underside of flagellomere, F2 slightly longer than wide (Fig. 54). T1 sparsely, but coarsely punctate. Membranous portion of

- gonostylus large, as in Fig. 12. Body length 8.0–10.0 mm
 ***S. kozlovi* Astafurova & Proshchalykin**
- 16 Tyloids on last flagellomeres (at least from F4 onward) usually cover more
 than 1/2 of ventral flagellar surfaces, often up to 4/5 Membranous portion of
 gonostylus larger, as in *S. kozlovi* (Fig. 12). Body length 7.0–11.0 mm
 ***S. pellucidus* Smith**
- Tyloids on last flagellomeres (at least from F4 onward) usually cover about 1/2
 of ventral flagellar surfaces, rarely up to 3/4. Membranous portion of gonosty-
 lus smaller (Fig. 6). Body length 6.0–9.0 mm ***S. ephippius* (Linné)**
- 17 Ocello-ocular area densely punctate, with punctures separated by about one
 puncture diameter. Gonostylus joining apex and partly inner surface of gono-
 coxite (Fig. 22). Body length 5.0–7.5 mm ***S. puncticeps* Thomson**
- Ocello-ocular area sparsely punctate, with punctures separated by 1–3 punc-
 ture diameters. Gonostylus joining only apex of gonocoxite (Fig. 8) **18**
- 18 F2 shorter, 1.4–1.6 times as long as wide. Tyloids on the flagellomeres extend
 across 1/4–1/2 of ventral flagellar surfaces. Body length 3.5–6.0 mm
 ***S. longulus* Hagens**
- F2 longer, 1.7–1.8 times as long as wide. Tyloids on the flagellomeres extend
 across 1/2–3/4 of ventral flagellar surfaces. Body length 4.0–5.0 mm
 ***S. turanicus* Astafurova & Proshchalykin**
- 19 T1 densely punctate. Face with appressed white pubescence below and above
 the antennal toruli **20**
- T1 impunctate or with sparse punctures (in *S. miniatus* sometimes relatively
 densely punctate). Face with appressed white pubescence only below the an-
 tennal toruli **22**
- 20 Tyloids variable, covering 1/2–4/5 flagellar ventral surfaces. Membranous
 portion of gonostylus small, triangular (Fig. 17)
 ***S. schwarzi* Astafurova & Proshchalykin**
- Tyloids covering from 3/4 to entire ventral flagellar surfaces. Membranous
 portion of gonostylus large, close to rectangular (Figs 10, 20) **21**
- 21 Antenna longer, F2 1.4 times as long as wide (Fig. 55). Membranous portion
 of gonostylus almost straight on inner edge (Fig. 20). Body length 5.0–7.5
 mm ***S. pinguiculus* Pérez**
- Antenna shorter, F2 1.2 times as long as wide (Fig. 56). Membranous por-
 tion of gonostylus weakly S-curved on inner edge (Fig. 10). Body length
 5.0–7.5 mm ***S. intermedius* Blüthgen**
- 22 Pronotum, between dorsal and lateral surfaces, rounded, not angulate
 (Fig. 36) **23**
- Pronotum, between dorsal and lateral surfaces, with sharp angle (Fig. 35) **26**

- 23 Tyloids on flagellomeres covering less than 1/3 of ventral flagellar surfaces. Membranous portion of gonostylus larger, trapezoidal (Fig. 5). Body length 6.0–9.0 mm.....***S. ferruginatus* Hagens**
- Tyloids on flagellomeres (at least from F4 onward) covering about 1/2–3/4 or entire of ventral flagellar surfaces. Membranous portion of gonostylus smaller, oval or almost square (Figs 13, 16, 24)**24**
- 24 Clypeus with fine, simple and sparsely plumose setae, sculpturing clearly visible (Fig. 33). Membranous portion of gonostylus square (Fig. 13). Body length 6.0–7.0 mm..... ***S. maruyamanus* Tsuneki**
- Clypeus with densely plumose setae, partly obscuring sculpturing (Fig. 34). Membranous portion of gonostylus close to oval**25**
- 25 Antenna short, middle flagellomeres as long as or slightly longer than wide. Tyloids on flagellomeres covering entire of ventral flagellar surfaces. Membranous portion of gonostylus longer, reach penis valve (Fig. 16). Body length 5.5–6.5 mm.....***S. murotai* Tsuneki**
- Antenna long, flagellomeres (from F3 onward) 1.2–1.3 times as long as wide. Tyloids on flagellomeres (at least from F4 onward) covering about 1/2–3/4 of ventral flagellar surfaces. Membranous portion of gonostylus shorter, not reach penis valve (Fig. 24). Body length 6.0–7.0 mm.....***S. tanoi* Tsuneki**
- 26 F2 short, 0.9–1.0 times as long as F3. Tyloids on flagellomeres (at least from F4 onward) usually cover entire ventral flagellar surfaces. Gonostylus with trapezoidal membranous portion (Fig. 7). Body length 5.0–6.5 mm.....
..... ***S. geoffrellus* (Kirby)**
- F2 longer, 1.1–1.2 as long as F3. Tyloids on flagellomeres shorter, covering at most 4/5 the ventral flagellar surfaces (in *S. miniatus* tyloids on last flagellomeres rare cover entire ventral flagellar surfaces).....**27**
- 27 Tyloids on flagellomeres covering more than 3/4 flagellar ventral surfaces. Gonostylus with large, trapezoidal membranous portion (Fig. 14). Body length 4.0–6.0 mm..... ***S. miniatus* Hagens**
- Felt-like areas on flagellomeres cover less 1/3 underside of flagellomere. Gonostylus with oval membranous portion or without one**28**
- 28 Head less transverse, 1.05 times as wide as long. Mesoscutum sparsely punctate, medially with punctures mostly separated by 1–3 puncture diameters. T1–T3 usually red, rarely terga entirely black. Gonostylus with oval membranous portion (Fig. 3). Body length 5.0–7.0 mm ***S. crassus* Thomson**
- Head more transverse, at least 1.15 times as wide as long. Mesoscutum very densely punctate, with confluent punctures (areolate). Terga usually wholly black, rare T1 dark red. Gonostylus without membranous portion (Fig. 11). Body length 7.5–8.5 mm.....***S. laticaudatus* Tsuneki**

Females

- 1 Hind wing with basal (*M*) vein weakly curved; costal margin with 7–14 hamuli. Usually large species: total body length 6.0–15.0 mm..... **2**
- Hind wing with basal (*M*) vein strongly curved; costal margin with 5–6 hamuli. Large or small species: total body length 5.5–11.0 mm..... **10**
- 2 Vertex less elevated (distance from top of head to upper margin of lateral ocellus less two lateral ocellar diameters as seen in frontal view), with longitudinal sharp carina (Fig. 37) **3**
- Vertex more elevated (distance from top of head to upper margin of lateral ocellus more two lateral ocellar diameters as seen in frontal view), acarinate, but sometimes with weak (indistinct) longitudinal ridge..... **5**
- 3 Face and gena with sparse, semi-erect, gray pubescence not obscuring integument. T1 with finer punctures (3–10 μm). Body length 6.0–8.0 mm ***S. cristatus* Hagens**
- Face and gena with dense, appressed, snow-white pubescence obscuring integument. T1 with coarser punctures (10–30 μm) **4**
- 4 Mesoscutum coarsely punctate (25–75 μm). T2 with marginal zone impunctate. Larger: body length 8.0–11.0 mm ***S. olivieri* Lepeletier de Saint Fargeau**
- Mesoscutum relatively finely punctate (25–40 μm). T2 with marginal zone distinctly punctate. Smaller: body length 6.5–8.5 mm ***S. pectoralis* Morawitz**
- 5 Gena flat. Preoccipital lateral carina developed (Fig. 40). Body length 9.0–12.0 mm ***S. scabricollis* Wesmael**
- Gena swollen. Preoccipital carina not developed **6**
- 6 Vertex shorter, distance from top of head to upper margin of lateral ocellus about 2 lateral ocellar diameters as seen in dorsal view (Fig. 42). T4 with marginal zone punctate and smooth between punctures or finely tessellate without punctures..... **7**
- Vertex longer, distance from top of head to upper margin of lateral ocellus equal to 2.5–3.0 lateral ocellar diameters as seen in dorsal view (Fig. 41). T4 with marginal zone impunctate, smooth (rarely indistinctly tessellate) **8**
- 7 T4 with marginal zone impunctate, finely tessellate (Fig. 46); T1 finely punctate (10–15 μm). Mesoscutum usually densely punctate, medially with punctures separated by not more than 1–3 puncture diameters, sometimes sparser. Body length 7.0–10.0 mm..... ***S. reticulatus* Thomson**
- T4 with marginal zone distinctly punctate, smooth between punctures (rarely indistinctly tessellate) (Fig. 45); T1 coarsely punctate (15–25 μm). Mesoscutum usually sparsely punctate, medially with punctures separated by mostly 2–4 puncture diameters. Body length 8.0–11.0 mm..... ***S. alternatus* Smith**
- 8 Mesoscutum densely punctate, with punctures separated by less than a puncture diameter (Fig. 47). Body length 9.0–12.0 mm ***S. albilabris* (Fabricius)**
- Mesoscutum sparsely punctate, medially with punctures separated by at least 2 puncture diameters (Fig. 50) **9**

- 9 Head rounded-rectangular on upper margin, square-shaped as seen in frontal view (Fig. 32); vertex sparsely punctate, punctures mostly separated by more than a puncture diameter. Pygidial plate equal or slightly narrower than metabasitarsus. T1 usually indistinctly punctate, with a few very fine punctures. Body length 7.0–10.0 mm..... ***S. monilicornis* (Kirby)**
- Head uniformly rounded on upper margin, oval as seen in frontal view (Fig. 28); vertex densely punctate, punctures mostly separated by less than a puncture diameter. Pygidial plate 0.5–0.6 times as wide as metabasitarsus. T1 distinctly punctate, with fine and coarser punctures. Body length 7.0–15.0 mm..... ***S. gibbus* (Linnaeus) and *S. nippon* Meyer***
- 10 Mandible simple (without an inner tooth) **11**
- Mandible bidentate..... **13**
- 11 Head narrower, at most 1.15 times as wide as long (Fig. 26). Body length 4.0–6.0 mm..... ***S. longulus* Hagens**
- Head broader, 1.2–1.3 times as wide as long..... **12**
- 12 Face, gena and mesepisternum with gray, sparse, semi-erect pubescence, not obscuring integument. Metasoma coarsely punctate (10–15 μ m). Pygidial plate as wide as metabasitarsus. Body length 5.0–8.0 mm.....
..... ***S. puncticeps* Thomson**
- Face, gena and mesepisternum with dense, snow-white, appressed, pubescence obscuring integument (Fig. 31). Metasoma finely punctate (3–5 μ m). Pygidial plate 1.2 times as wide as metabasitarsus
..... ***S. turanicus* Astafurova & Proshchalykin**
- 13 Pygidial plate at least 1.2 times wider than metabasitarsus, usually dull. Mesoscutum densely punctate, punctures usually separated by less than two puncture diameters. Total body length 7.0–11.0 mm..... **14**
- Pygidial plate equal to or narrower than metabasitarsus, shiny. Mesoscutum usually sparsely punctate, disc medially with punctures separated by more than two puncture diameters. Total body length 4.0–9.0 mm..... **18**
- 14 Vertex with longitudinal carina (Figs 38, 39) **15**
- Vertex without longitudinal carina..... **16**
- 15 Vertex with obvious longitudinal carina (Fig. 38). Setae on scape shorter than width of scape. Pygidial plate 1.2–1.4 times wider than metabasitarsus. Body length 7.0–9.0 mm..... ***S. pieli* Cockerell**
- Vertex with weakly visible longitudinal carina (Fig. 39). Setae on scape distinctly longer than width of scape (not obviously in old females). Pygidial plate 1.4–1.5 times wider than metabasitarsus. Body length 8.0–9.0 mm.....
..... ***S. kozlovi* Astafurova & Proshchalykin**
- 16 Pygidial plate 1.6–1.7 times as wide as metabasitarsus. Gena wider, 0.8 times as wide as eye in lateral view. Mesoscutum densely punctate, with punctures

* Females of this vicarious species are very difficult to distinguish morphologically; however, *S. nippon* is distributed in China to Gansu on the West, whereas *S. gibbus* is distributed in China to Xinjiang on the East.

- separated by at most a puncture diameter (Fig. 51). Body length 7.5–10.0 mm.....***S. laticaudatus* Tsuneki**
- Pygidial plate 1.2–1.5 times as wide as metabasitarsus. Gena narrower, 0.7 times as wide as eye in lateral view. Mesoscutum sparsely punctate, medially with punctures usually separated by 1–2 puncture diameters (Fig. 52).....**17**
- 17 Head more transverse, 1.30–1.35 times as wide as long; vertex, behind ocelli, not elevated in frontal view. Setae on scape distinctly longer than width of scape. Pygidial plate 1.3–1.5 times as wide as metabasitarsus..... ***S. pellucidus* Smith**
- Head less transverse, 1.20–1.25 times wider than long; vertex, behind ocelli, weakly elevated. Setae on scape shorter than width of scape. Pygidial plate 1.2–1.4 times as wide as metabasitarsus ***S. ephippius* (Linné) and *S. grahami* Cockerell***
- 18 Clypeus densely punctate, punctures separated by less than one puncture diameter. Pronotum, between dorsal and lateral surfaces, rounded, not angulate (Fig. 36).....**19**
- Clypeus sparsely punctate, punctures separated by at least one puncture diameter. Pronotum, between dorsal and lateral surfaces, with sharp angle (Fig. 35).....**22**
- 19 Hind femur narrow, regularly pointed toward distal end, its length more than 3.5 times its maximum width. Body length 6.0–7.5 mm..... ***S. maruyamanus* Tsuneki**
- Hind femur widened in proximal half, its length at most 3 times its maximum width.....**20**
- 20 Vertex, behind ocelli, weakly elevated in frontal view (Fig. 25). Body length 6.0–9.0 mm.....***S. ferruginatus* Hagens**
- Vertex not elevated in frontal view (Fig. 30).....**21**
- 21 Thorax ventrally with sculpture finer than on sides (Fig. 44). Pygidial plate slightly narrower than metabasitarsus. Body length 6–7 mm ***S. tanoi* Tsuneki**
- Thorax ventrally with sculpture as coarse as that on sides (Fig. 43). Pygidial plate as wide as metabasitarsus. Body length 5.5–6.5 mm..... ***S. murotai* Tsuneki**
- 22 Vertex longer, distance from top of head to upper margin of lateral ocellus equal to about 3–3.5 times lateral ocellar diameters as seen in dorsal view. Upper half of gena with appressed, dense pubescence obscuring integument..... **23**
- Vertex shorter, distance from top of head to upper margin of lateral ocellus equal to about two lateral ocellar diameters as seen in dorsal view. Gena with erect, sparse pubescence**24**

** Females of these species are very difficult to distinguish morphologically; however, *S. ephippius* is distributed in North-West China (Xinjiang), whereas *S. grahami* is recorded from Central, South and East China; the male of *S. grahami* is unknown.

- 23 Mesoscutum and mesoscutellum very sparsely punctate, with tiny punctures separated by 1–7 diameters (Fig. 48). Body length 5.0–7.0 mm *S. pinguiculus* Pérez
- Mesoscutum and mesoscutellum more densely punctate, with coarse punctures separated by 1–3 puncture diameters (Fig. 49). Body length 6.5–8.5 mm..... *S. intermedius* Blüthgen
- 24 F3 transverse, 0.6–0.7 times as long as wide, as long as F1. Pygidial plate 0.9–1.0 as wide as metabasitarsus.....25
- F3 square, as long as wide, longer than F1 Pygidial plate 0.6–0.8 as wide as metabasitarsus.....26
- 25 Paraocular area with dense, strongly plumose setae below the antennal toruli (Fig. 29). Body length 4.5–5.5 mm *S. schwarzi* Astafurova & Proshchalykin
- Face with sparse, simple and weakly plumose setae (Fig. 27). Body length 4.0–6.0 mm..... *S. miniatus* Hagens
- 26 Head more transverse, 1.25 times as wide as long. Labrum trapezoidal, 0.7 times as long as wide. Hind femur strongly enlarged on proximal half, maximum width 0.4 times its length. Body length 5.0–8.0 mm *S. crassus* Thomson
- Head less transverse, 1.1 times as wide as long. Labrum semicircular, 0.5 times as long as width. Hind femur weakly enlarged on proximal half, maximum width 0.35 times its length. Body length 4.5–6.5 mm..... *S. geoffrellus* (Kirby)


List of species

Sphecodes albilabris (Fabricius, 1793)

Figs 1, 47

Material examined. CHINA: *Liaoning*, 1 ♀, 50 km N Mukden [Shenyang] [42°12'N, 123°23'E], 20.VII.1952, leg. Rubtsov (ZISP); *Inner Mongolia*, 1 ♂, Xilinhot [43°54'N, 116°00'E], 27.VII.1971, leg. Y.-W. Zhang (IZCAS); *Hebei*, 1 ♀, Kreis Yongnian [36°25'N, 114°14'E], 1995, leg. S.-Q. Li (IZCAS); 1 ♀, Yangjiaping [39°58'N, 115°24'E], 3.VIII.1937, leg. O. Piel (IZCAS); 1 ♀, Yu Xian, Xihey-ing [39°57'N, 114°00'E], 800 m, 29.VII.1964, leg. B.-Q. Li (IZCAS); *Beijing*, 1 ♀, Xiyuan [39°55'N, 116°24'E], 50 m, 23.VII.1962, leg. C.-G. Wang (IZCAS); *Shanxi*, 1 ♀, Xiexian, Zhongtiao Shan Mts. [34°48'N, 111°36'E], 22–24.V.1996, leg. JH (PCMS); *Gansu*, 1 ♂, Lanzhou [36°00'N, 103°25'E], 1500 m, 9.IX.1957, leg. Y.-R. Zhang (IZCAS).

Distribution. *China (Liaoning, Inner Mongolia, Hebei, Beijing, Shanxi, Gansu), Central Asia, Russia, Europe (north to Finland and Sweden), Turkey, Syria, Caucasus, North Africa, Israel, India.



Figures 1–12. Genitalia, males, dorsal view. **1** *Sphecodes albilabris* (Fabricius) **2** *S. alternatus* Smith **3** *S. crassus* Thomson **4** *S. cristatus* Hagens **5** *S. ferruginatus* Hagens **6** *S. ephippius* (Linné) **7** *S. geoffrellus* (Kirby) **8** *S. longulus* Hagens **9** *S. gibbus* (Linnaeus) **10** *S. intermedius* Blüthgen **11** *S. laticaudatus* Tsuneki **12** *S. kozlovi* Astafurova & Proshchalykin. Scale bars: 0.25 mm.

***Sphecodes alternatus* Smith, 1853**

Figs 2, 46

Material examined. CHINA: *Gansu*, 2 ♂♂, oasis Sachjou [Dunhuang] [40°09'N, 94°40'E], Gashun Gobi desert, 1–3.VIII.1895, VR, PK (ZISP); *Xinjiang*, 1 ♂, Yin-
ing, Boro Hqro Mts [44°06'N, 81°00'E], 27.VII.1991, Snizek (PCMS); 1 ♀, 37 ♂♂,
Bugas near Khami [43°14'N, 93°50'E], 20.VIII–8.IX.1895, VR, PK (ZISP).

Distribution. *China (Gansu, Xinjiang), Central Asia, Europe, Russia (south of
European part and east to Khakassia Republic), Turkey, Iran, North Africa.

***Sphecodes crassus* Thomson, 1870**

Figure 3

Material examined. CHINA: *Inner Mongolia*, 2 ♀♀, Suburgan-gol, Alashan [Helan
Shan] Mt., Gobi, 29–30.VI.1908, PK (ZISP); 6 ♀♀, Tszosto, Alashan [Helan Shan]
Mt., Gobi, 13–14.V.1908, PK (ZISP); *Shanxi*, 1 ♀, Xiexian, Zhongtiao Shan Mts
[34°48'N, 111°36'E], 22–24.V.1996, leg. JH (PCMS); 2 ♀♀, 13 km S Yichuan
[35°54'N, 110°36'E], 19.V.1996, leg. JH (PCMS).

Distribution. *China (Inner Mongolia, Shanxi), Central Asia, Mongolia, Russia,
Europe (north to 64°), Caucasus, Turkey, Iran, Japan, North Africa.

***Sphecodes cristatus* Hagens, 1882**

Figure 4

Sphecodes alfkeni Meyer, 1922: 172, ♀ (lectotype: ♀, **designated here**, China, Tientsin
[Tianjin], [leg.] Weber, MNHB, examined). – Syn. n.

Material examined. CHINA: *Heilongjiang*, 1 ♀, Harbin [45°46'N, 126°39'E], 8.X.1952
(IZCAS); 1 ♀, idem, 27.VII.1952 (IZCAS); 1 ♀, idem, 19.VII.1953 (IZCAS); 1 ♀,
idem, 25.VII.1955 (IZCAS); 1 ♂, idem, 8.X.1952 (IZCAS); 1 ♂, idem, 16.VII.1952
(IZCAS); *Jilin*, 1 ♀, Gongzhuling [43°79'N, 124°69'E], 9.VI.1962, leg. T.-L. Cheng (IZ-
CAS); *Liaoning*, 1 ♂, 50 km N Mukden [Shenyang], 20.VII.1952, Rubtsov (ZISP); 1 ♀,
Guicheng [43°40'N, 126°15'E], 30.VII.1962, leg. T.-L. Cheng (IZCAS); *Inner Mongolia*,
4 ♂♂, 2 ♀♀, Dingyuanying [Bayan Hot], Alashan [Helan Shan] Mt., 22.V., 26.V., 17–
26.IX.1908, PK (ZISP); 1 ♂, Tszosto, Alashan [Helan Shan] Mt., Gobi, 13–14.V.1908,
PK (ZISP); 1 ♂, Ulanqab Men, Tomortei [41°48'N, 113°06'E], 31.VIII.1971 (IZCAS);
2 ♂♂, Hailar Shi [49°12'N, 119°42'E], 3.VIII.2006, leg. H.-Y. Zhang (IZCAS), 2 ♂♂,
idem, 2.VIII.2006, leg. Y.-J. Li (IZCAS), 1 ♂, idem, 8.VIII.2006, leg. P. Wang (IZCAS); 2
♂♂, Ordos, Dundatu [37°43'N, 108°10'E], 24.VII.2006, leg. Y.-J. Li (IZCAS); 2 ♀♀, 1
♂, idem, 25.VII.2006, leg. P. Wang (IZCAS), 5 ♀♀, 3 ♂♂, idem, 25.VII.2006, leg. H.-Y.
Zhang (IZCAS); 3 ♀♀, idem, 25.VII.2006, leg. M. Luo (IZCAS); 1 ♀, 1 ♂, Hohhot Shi,

Heling Xian, Mengniu Zheng [40°49'N, 111°39'E], 15.VII.2006, leg. Y.-J. Li (IZCAS); 1 ♂, Hulun Buir Meng, Manzhouli Shi [49°12'N, 119°45'E], 5.VIII.2006, leg. Y.-J. Li (IZCAS); 1 ♀, Uxin Qi, Batugou [38°38'N, 108°53'E], 28.VII.2006, leg. M. Luo (IZCAS); *Hebei*, 1 ♀, Yangyuan Xian, Liufang [40°11'N, 114°28'E], 950 m, 12.IV.2002, leg. Z.-Q. Niu (IZCAS); *Tianjin*, 1 ♀, Balitai [38°57'N, 117°19'E], 13.X.1953, leg. Z.-R. Yu (IZCAS). *Beijing*, 1 ♀, 1 ♂, Xiangshan [39°54'N, 116°12'E], 100 m, 19.IX.1962, leg. Y.-S. Shi (IZCAS); 3 ♂♂, Beijing [39°55'N, 116°24'E], 28.VIII.1973, leg. S.-F. Wang (IZCAS); 1 ♀, Wofosi [40°03'N, 115°10'E], 100 m, 10.V.1962, leg. S.-Y. Wang (IZCAS); 1 ♂, idem, 18.IX.1981, leg. Q. Zhou (IZCAS); 1 ♀, Zizhuyuan [39°57'N, 116°19'E], 24.IV.1962, leg. S.-M. Ge (IZCAS); *Shanxi*, 1 ♀, Qingjian env. [36°54'N, 110°36'E], 15.V.1996, leg. JH (PCMS); 1 ♀, Monan [34°42'N, 111°42'E], 26–28.V.1996, leg. JH (PCMS); 1 ♀, Suide, [37°18'N, 110°42'E], 13–14.V.1996, leg. JH (PCMS); *Shandong*, 1 ♀, Jinan [36°48'N, 117°01'E], 24.VI.1937 (IZCAS); *Shaanxi*, 1 ♀, Gangui [36°48'N, 110°18'E], 35 km NE Yanan, 17–18.V.1996, leg. JH (PCMS); *Ningxia*, 1 ♂, Ningxia [Yinchuan], Ordos, Gobi, 1–4.VI.1908, PK (ZISP); 6 ♂♂, Yanchi Xian, Sidunzi [37°28'N, 107°09'E], 1455 m, 22.VI.2016, leg. Z.-Q. Niu, D. Zhang (IZCAS); *Xinjiang*, 1 ♀, Jimsar Xian [44°00'N 89°03'E], 14.V.1955, leg. S.-J. Ma, K.-L. Xia, Y.-L. Cheng (IZCAS); 1 ♀, Jeminay Xian, S229, 14 km [47°14'N, 85°19'E], 1080 m, 28.VIII.2002, leg. Z.-Q. Niu (IZCAS); 1 ♀, Tian Shan [43°10'N, 86°00'E], 28.VIII.1957, leg. G. Wang (IZCAS); 1 ♀, Manas Xian, Shihezi [44°07'N, 86°00'E], 550 m, 6.VI.1957, leg. C.-P. Hong (IZCAS); 1 ♀, idem, 590 m, 27.VIII.1957, leg. S.-Y. Wang (IZCAS); 1 ♀, Yining Xian [44°00'N, 81°21'E], 4.VIII.1957, leg. W.-Y. Yang (IZCAS).

Published records. Blüthgen 1927: 42, as *Sphecodes alfkeni* Meyer (Tianjin); Ascher and Pickering 2018 (Beijing).

Distribution. China (*Heilongjiang, *Jilin, *Liaoning, *Inner Mongolia, *Hebei, Tianjin, Beijing, *Shanxi, *Shandong, *Shaanxi, *Ningxia, *Xinjiang), Europe (north to Sweden), Korea, Russia, Caucasus, Turkey, Central Asia, Mongolia.

Sphecodes ephippius (Linné, 1767)

Figure 6

Material examined. CHINA: *Xinjiang*, 1 ♂, Yaerkate [42°52'N, 92°50'E], 3.VIII.1956, leg. W.-Y. Yang (IZCAS).

Distribution. *China (Xinjiang), Russia (east to Irkutsk Prov.), Mongolia, Central Asia, Caucasus, Turkey, Europe (north to 62°).

Sphecodes ferruginatus Hagens, 1882

Figs 5, 25, 36

Material examined. CHINA: *Beijing*, 2 ♀♀, 3 ♀♀, Bada Ling, Sanbu [40°22'N, 115°58'E], 500 m, 18, 27.IV.2002, leg. Z.-Q. Niu (IZCAS); 1 ♀, Miaofengshan

[40°01'N, 115°59'E], 24.V.1957, leg. M.-H. Wang (IZCAS); 1 ♀, Wofosi [40°03'N, 115°10'E], 15.V.1961, leg. S.-M. Ge (IZCAS); *Shanxi*, 1 ♀, Xiexian [34°48'N, 111°36'E], Zhongtiao Shan Mts., 22–24.V.1996, leg. JH (PCMS).

Distribution. *China (Beijing, Shanxi), Central Asia, Russia, Europe (north to 66°), Caucasus, Turkey, Japan.

Sphecodes geoffrellus (Kirby 1802)

Figure 7

Material examined. CHINA: *Heilongjiang*, 1 ♂, Da Hinggan Ling [51°42'N, 126°36'E], 23.VII.1980 (IZCAS); *Shaanxi*, 11 ♀♀, Gangui [36°48'N, 110°18'E], 35 km NE Yanan, 17–18.V.1996, leg. JH (PCMS); 1 ♀, 13 km S Yichuan [35°54'N, 110°36'E], 19.V.1996, leg. JH (PCMS).

Distribution. *China (Heilongjiang, Shanxi), Central Asia, Europe (north to 66°), Russia (east to Far East), Turkey, Near East, Mongolia, Japan.

Sphecodes gibbus (Linnaeus, 1758)

Figs 9, 41, 57

Sphecodes gibbus var. *turkestanicus* Meyer, 1920: 113 (holotype: 1 ♀, Uzbekistan: Golodnaja Steppe [Gulistan], MNHB). Synonymized by Blüthgen 1923: 510.

Material examined. CHINA: *Xinjiang*, 1 ♀, 13 ♂♂, Bugas near Khami [43°14'N, 93°50'E], 20.VIII–8.IX.1895, leg. VR, PK (ZISP); 1 ♂, Qitai Xian [44°31'N, 90°06'E], 10.VII.1975 (IZCAS); 1 ♀, Kashi [39°14'N, 75°32'E], 133 m, 10.VII.1959, leg. C.-Q. Li (IZCAS); 1 ♂, idem, 10.VII.1959, leg. A-F. Tian (IZCAS).

Published records. Meyer, 1920: Yarkand (Xinjiang), as *Sphecodes gibbus* var. *turkestanicus* Meyer, 1920.


Distribution. China (Xinjiang), Central Asia, Russia (east to Yakutia), Pakistan, Mongolia, Europe (north to 63°), Turkey, Israel, North Africa, India.

Sphecodes grahami Cockerell, 1922

Figure 52

Sphecodes grahami Cockerell, 1922: 12 (holotype: ♀, China, Sichuan: Suifu, Graham coll.; USNM);

Material examined. CHINA: *Shanxi*, 1 ♀, Xiexian [34°48'N, 111°36'E], Zhongtiao Shan Mts., 22–24.V.1996, leg. JH (PCMS); *Shaanxi*, 2 ♀♀, Gangui [36°48'N, 110°18'E], 35 km NE Yanan, 17–18.V.1996, leg. JH (PCMS).



Figures 13–24. Genitalia, males, dorsal view. **13** *Sphecodes maruyamanus* Tsuneki **14** *S. miniatus* Hagens **15** *S. monilicornis* (Kirby) **16** *S. murotai* Tsuneki **17** *S. schwarzi* Astafurova & Proshchalykin **18** *S. nurekensis* Warncke **19** *S. pieli* Cockerell **20** *S. pinguiculus* Pérez **21** *S. olivieri* Lepeletier de Saint Fargeau **22** *S. puncticeps* Thomson **23** *S. scabricollis* Wesmael **24** *S. tanoi* Tsuneki. Scale bars: 0.25 mm.

Published records. Cockerell 1922: 12 (Shanghai); Ascher and Pickering 2018 (Jilin, Hebei, Anhui, Jiangsu, Shanghai, Zhejiang, Yunnan, Xizang, Guandong).

Distribution. China (Jilin, Hebei, *Shanxi, *Shaanxi, Anhui, Jiangsu, Shanghai, Shanghai, Zhejiang, Sichuan, Yunnan, Xizang, Guandong).

Remark. The female of this species is challenging to distinguish from West-Palaearctic *S. ephippius* (Linné, 1767).

***Sphecodes intermedius* Blüthgen, 1923**

Figs 10, 49, 56

Material examined. CHINA: *Gansu*, 1 ♂, Shibendu, oasis Sachjou [Dunhuang] [40°09'N, 94°40'E], Gashun Gobi desert, 9–12.VIII.1895, leg. VR, PK (ZISP).

Distribution. *China (Gansu), Central Asia, Europe, Russia (European part, Ural), North Africa, Caucasus, Turkey.

***Sphecodes kozlovi* Astafurova & Proshchalykin, 2015**

Figs 12, 39, 54

Material examined. CHINA: *Inner Mongolia*, 3 ♀♀, Tszosto, Alashan [Helan Shan] Mt., Gobi, 10–18.V.1908, leg. PK (ZISP); 7 ♀♀, Dingyuanying [Bayan Hot], Alashan [Helan Shan] Mt., 10, 18–19.VI.1908, 15–16. IV.1909, leg. PK (ZISP); *Shanxi*, 1 ♀, Monan [34°42'N, 111°42'E], 26–28.V.1996, leg. JH (PCMS); *Ningxia*, 2 ♀♀, Yanchi [37°24'N, 107°36'E], 11.V.1996, leg. JH (PCMS).

Distribution. *China (Inner Mongolia, Shanxi, Ningxia), Mongolia (Dornod Aimag, Khentii Aimag).

***Sphecodes laticaudatus* Tsuneki, 1983**

Figs 11, 51

Material examined. CHINA: *Hebei*, 1 ♂, Xinglong Xian, Wuling Shan [40°26'N, 117°31'E], 28.VIII.1973 (IZCAS).


Distribution. *China (Hebei), Russia (Far East), Japan.

***Sphecodes longulus* Hagens, 1882**

Figs 8, 26

Sphecodes subfasciatus Blüthgen, 1934: 22, ♀ (holotype: ♀, China, S. Kansu, 19.VI.1930, leg. Hummel, NHRS, examined). – Syn. n.

Material examined. CHINA: *Inner Mongolia*, 1 ♀, Goytzo valley, Alashan, Gobi, 5.IV.1908, leg. PK (ZISP); *Hebei*, 1 ♀, Changli Xian [39°38'N, 119°05'E], 28.IV.1962, leg. T.-L. Cheng (IZCAS); 1 ♀, Xiaowutai Shan [38°36'N, 115°39'E], 1200 m, 22.VIII.1964, leg. Y.-H. Han (IZCAS); *Shaanxi*, 1 ♀, Gangui [36°48'N,



Figures 25–30. Head, females, frontal view. **25** *Sphecodes ferruginatus* Hagens **26** *S. longulus* Hagens **27** *S. miniatus* Hagens **28** *S. nippon* Meyer **29** *S. schwarzi* Astafurova & Proshchalykin **30** *S. tanoi* Tsuneki. Scale bars: 0.5 mm.

110°18'E], 35 km NE Yanan, 17–18.V.1996, leg. JH (PCMS); *Gansu*, 1 ♀, Lanzhou [36°00'N, 103°25'E], 27.IV.1955, leg. S.-J. Ma, K.-L. Xia, Y.-L. Cheng (IZ-CAS); *Xinjiang*, 1 ♂, Tacheng Xian [46°25'N, 82°32'E], 470 m, 10.IX.1960, leg.

S.-Y. Wang (IZCAS); 1 ♂, Bostanterak [39°07'N, 95°03'E], 9.VII.1959, leg. S.-Y. Wang (IZCAS).

Distribution. China (*Inner Mongolia, *Hebei, *Shaanxi, Gansu, *Xinjiang), Central Asia, Russia, Europe (north to Finland, Sweden, Denmark, England), Turkey, Syria, Japan.

Sphecodes manchurianus Strand & Yasumatsu, 1938

Sphecodes manchurianus Strand & Yasumatsu, 1938: 80 (holotype: ♂, China : “Fengtien (Mukden), South Manchoukuo, 5.VIII.1930, T. Nozawa leg.”; KUF, lost).

Material examined. No material examined.

Distribution. China (Liaoning).

Remark. Known only from the holotype.

Sphecodes monilicornis (Kirby, 1802)

Figs 15, 32

Material examined. CHINA: *Heilongjiang*, 1 ♂, Morin Dawa [47°21'N, 128°03'E], 24.VII.1976 (IZCAS); 1 ♂, Harbin [45°46'N, 126°39'E], 6.VII.1947 (IZCAS); 1 ♀, idem, 27.VII.1952 (IZCAS); 2 ♂♂, idem, 25.VII.1953 (IZCAS); 7 ♂♂, idem, 4.VII.1955 (IZCAS); 1 ♂, idem, 8.VII.1955 (IZCAS); 1 ♀, 3 ♂♂, idem, 10.VII.1955 (IZCAS); 1 ♀, 5 ♂♂, idem, 19.VII.1955 (IZCAS); 5 ♂♂, idem, 10.VII.1955 (IZCAS); 1 ♂, idem, 11.VII.1955 (IZCAS); 3 ♂♂, idem, 23.VII.1953 (IZCAS); 1 ♂, idem, 9.VIII.1955 (IZCAS).


Distribution. *China (Heilongjiang), Central Asia, Mongolia, Russia, North Pakistan, Europe (north to 64°), Caucasus, Turkey, North Africa.

Sphecodes nippon Meyer, 1922


Figs 28, 50, 58

Sphecodes kansuensis Blüthgen, 1934: 21, fig. 11, ♂ (holotype: ♂, China, S. Kansu [Gansu] 19.VI.1930, leg. Hummel, NHRS, examined). – Syn. n.

Material examined. CHINA: *Heilongjiang*, 1 ♂, Harbin [45°46'N, 126°39'E], 24.IX.1950; 1 ♂, idem, 16.VII.1952; 1 ♂, idem, 25.VII.1952; 1 ♂, idem, 23.VII.1953; 1 ♂, idem, 11.VII.1954; 1 ♂, idem, 4.VII.1955; 4 ♂♂, idem, 8.VII.1955; 2 ♂♂, idem, 25.VII.1955 (IZCAS); 2 ♂♂, Mao'ershan [47°21'N, 128°03'E], 29.VII.1951 (IZCAS); 1 ♂, Hengdaohezi [45°57'N, 129°57'E], 28.VII.1951 (IZCAS); *Inner Mongolia*, 1 ♂, 3 ♀♀, Dingyuanying [Bayan Hot], Alashan [Helan Shan] Mt., 16–17.V., 3–6.VI., 11–16.IX.1908, PK (ZISP); *Hebei*, 1 ♂, Yangjiaping [39°58'N,




Figures 31–36. Diagnostic characters of *Sphecodes* species. **31, 32, 35, 36** Females **33, 34** Males **31–34** head, frontal view **35, 36** pronotum, lateral, view **31** *Sphecodes turanicus* Astafurova & Proshchalykin **32** *S. monilicornis* (Kirby) **33** *S. maruyamanus* Tsuneki **34** *S. murotai* Tsuneki **35** *S. pellucidus* Smith **36** *S. ferruginatus* Hagens. Scale bars: 0.5 mm.



Figures 37–40. Diagnostic characters of *Sphecodes* species, females. **37–39** vertex, dorso-lateral view **40** preoccipital carina and gena, dorso-lateral view **37** *Sphecodes olivieri* Lepeletier de Saint Fargeau **38** *S. pieli* Cockerell **39** *S. kozlovi* Astafurova & Proshchalykin **40** *S. scabricollis* Wesmael. Scale bars: 0.25 mm.

115°24'E], 17.VII.1937; 1 ♂, idem, 20.VII.1937; 1 ♀, idem, 6.VII.1937; 1 ♀, idem, 8.VII.1937, leg. O. Piel (IZCAS); 1 ♂, Xiaowutai Shan [38°36'N, 115°39'E], 1200 m, 25.VIII.1964; 1 ♂, idem, 11.VII.1964; 1 ♂, idem, 12.VII.1964, leg. Y.-H. Han (IZCAS); 1 ♂, Xinglong Xian, Wuling Shan [40°26'N, 117°31'E], 28.VIII.1973 (IZCAS); *Tianjin*, 1 ♀, Balitai [38°57'N, 117°19'E], 24.IV.1953, leg. Z.-Y. Xu (IZCAS); *Beijing*, 1 ♂, 40 km N Beijing [40°09'N, 116°14'E], 28.IX.1952, Rubtsov (ZISP); 1 ♀, Bada Ling [40°22'N, 115°58'E], 700 m, 2.VII.1974, leg. Y.-S. Shi (IZCAS); 1 ♀, Bada Ling, Sanbu [40°22'N, 115°58'E], 500 m, 27.IV.2002, leg. Z.-Q. Niu (IZCAS); 1 ♀, Xidazhuangke village, Songshan [40°31'N, 115°47'E], 910 m, 15.V.2007, leg. H.-R. Huang (IZCAS); 1 ♀, Miaofengshan [40°01'N, 115°59'E], 2.VIII.1957; 1 ♂, idem, 18.VII.1957, leg. M.-H. Wang (IZCAS); *Shaanxi*, 9 ♀♀, Gangui [36°48'N, 110°18'E], 35 km NE Yanan, 17–18.V.1996, leg. JH (PCMS); *Gansu*, 5 ♂♂, oasis Sachjou [Dunhuang] [40°09'N, 94°40'E], Gashun Gobi desert, 28–30.VII.1895, leg. VR, PK; 1 ♀, Lanzhou, 25.VII.1908, leg. PK (ZISP).



Figures 41–46. Diagnostic characters of *Sphecoides* species, females. **41, 42** head, dorsal view **43, 44** thorax, ventral view **45, 46** T4-T5, dorsal view **41** *S. gibbus* (Linnaeus) **42, 45** *S. reticulatus* Thomson **43** *S. murotai* Tsuneki **44** *S. tanoi* Tsuneki **46** *S. alternatus* Smith. Scale bars: 0.5 mm.


Distribution. China (*Heilongjiang, *Inner Mongolia, *Hebei, *Tianjin, *Beijing, *Shaanxi, Gansu), Russia (East Siberia, Far East), Mongolia, Japan.

***Sphecoides nurekensis* Warncke, 1992**

Figure 18

Material examined. CHINA: *Xinjiang*, 1 ♂, Bugas near Khami [N43°14' E93°50'], 20.VIII.1895, leg. VR, PK (ZISP); 1 ♂, Ürümqi [43°28'N, 87°32'E], 980 m, 2.IX.1959, leg. S.-Y. Wang (IZCAS).

Distribution. *China (Xinjiang), Tajikistan.



Figures 47–52. Scutum, females, dorsal view. **47** *S. albilabris* (Fabricius) **48** *S. pinguiculus* Pérez **49** *S. intermedius* Blüthgen **50** *S. nippon* Meyer **51** *S. laticaudatus* Tsuneki **52** *S. grahami* Cockerell. Scale bars: 0.5 mm.

***Sphecodes olivieri* Lepeletier de Saint-Fargeau, 1825**

Figs 21, 37

Material examined. CHINA: *Gansu*, 1 ♀, 2 ♂♂, oasis Sachjou [Dunhuang] [40°09'N, 94°40'E], Gashun Gobi desert, 24.VII, 1–3.VIII.1895, leg. VR, PK (ZISP); 1 ♂, Zhangye [38°32'N, 100°14'E], 1450 m, 29.VII.1957, leg. Y.-R. Zhang (IZCAS); *Xinjiang*, 117 ♂♂, Bugas near Khami [43°14'N, 93°50'E], 20.VIII–8.IX.1895, leg.

VR, PK (ZISP); 3 ♀♀, Manas Xian [44°10'N, 86°07'E], 400 m, 9.VI.1953, leg. C.-P. Hong (IZCAS); 1 ♀, idem, 9.VI.1953, leg. W.-Y. Yang (IZCAS); 1 ♂, Manas Xian, Shihezi [44°07'N, 86°00'E], 500 m, 27.VIII.1959, leg. C.-Q. Li (IZCAS); 1 ♂, Burqin Xian [47°25'N, 86°32'E], 480 m, 25.VIII.1960, leg. S.-Y. Wang (IZCAS); 1 ♂, Turpan Xian [42°32'N, 89°07'E], 30.VI.1958 (IZCAS).

Distribution. *China (Gansu, Xinjiang), Central Asia, South Europe, Russia (south of European part), Caucasus, Turkey, Iran, Pakistan, India, Israel, United Arab Emirates, Qatar, North Africa.

Sphecodes pectoralis Morawitz, 1876

Material examined. CHINA: *Gansu*, 1 ♀, 1 ♂, oasis Sachjou [Dunhuang] [40°09'N, 94°40'E], Gashun Gobi desert, 28.VII–4.VIII.1895, leg. VR, PK (ZISP); *Xinjiang*: 4 ♀♀, 140 ♂, Bugas near Khami [43°14'N, 93°50'E], 20.VIII–8.IX.1895, leg. VR, PK (ZISP).

Distribution. *China (Gansu, Xinjiang), Central Asia.

Sphecodes pellucidus Smith, 1845

Figure 35

Sphecodes pellucidus var. *hybridus* Blüthgen, 1924: 516, ♀ (syntypes: ♀♀, China: Sichuan, NHRS). Synonymized by Warncke 1992: 20.

Sphecodes pellucidus var. *niveipennis* Meyer, 1925: 7, ♂ (lectotype: ♂, **designated here**, Chin. Turkestan, Uss-Lusch. Jarkand [China, Xinjiang, Yarkand] 1600 m, 4–6.8.90, Conrandt S. / *pellucidus* v. *niveipennis* Dr. R Meyer det.; MNHB, examined). Synonymized by Warncke 1992: 20.

Material examined. CHINA: *Gansu*, 1 ♀, Lanzhou, 11–25.III.1901, leg. PK (ZISP); 3 ♂♂, Dankhe River, S to Sachzhou [Dunhuang], Gashunskoe Gobi [39°55'N, 94°20'E], 24.VII.1895, leg. VR, PK (ZISP).

Distribution. China (*Gansu, Xinjiang, Sichuan), Central Asia, Russia, Mongolia, Europe (north to 66°), Caucasus, Turkey, North Africa.

Sphecodes pieli Cockerell, 1931

Figs 19, 38, 53

Sphecodes pieli Cockerell, 1931: 13, ♂ (holotype: ♂, China, Shanghai, Zo-Se, June 16, 1930 (Piel No 34), USNM).

Sphecodes orientalis Astafurova & Proshchalykin, 2014: 517–518, ♀, ♂ (holotype: ♂, Russia, Primorskiy Terr.: 15 km SW Slavyanka, 31.VIII.1995, S. Belokobylskij, ZISP, examined). – Syn. n.

Material examined. CHINA: *Hebei*, 1 ♀, Xiaowutai Shan [38°36'N, 115°39'E], 1200 m, 19.VI.1964, leg. Y.-H. Han (IZCAS); *Beijing*, 1 ♀, Bada Ling, Sanbu [40°22'N, 115°58'E], 500 m, 27.IV.2002 (IZCAS); 2 ♀♀, idem, 28.IV.2002, leg. Z.-Q. Niu (IZCAS); 1 ♀, Xidazhuangke village, Songshan [40°31'N, 115°47'E], 910 m, 15.V.2007, leg. H.-R. Huang (IZCAS); 1 ♀, Mentougou, Xiaolongmen, Liyuanling [39°58'N, 115°28'E], 1140–1250 m, 19.V.2002, leg. Z.-Q. Niu (IZCAS); *Shaanxi*, 1 ♀, Gangui, 35 km NE Yanan [36°48'N, 110°18'E], 17–18.V.1996, leg. JH (PCMS); 1 ♀, 13 km S Yichuan [35°59'N, 110°36'E], 19.V.1996, leg. JH (PCMS); 1 ♀, Jingangling, 50 km W Linfen, [36°12'N, 111°42'E], 29–30.V.1996, leg. JH (PCMS); *Sichuan*, 1 ♀, Nanping, Ta Zang [33°15'N, 104°15'E], 2200 m, 15–18.VI.1990, JH (PCMS).

Published records. Ascher and Pickering, 2018 (Zhejiang, Jiangsu).

Distribution. China (*Hebei, *Beijing, *Shaanxi, Jiangsu, Shanghai, Zhejiang, *Sichuan), Russia (Far East).

Sphecodes pinguiculus Pérez, 1903

Figs 20, 48, 55


Material examined. CHINA: *Inner Mongolia*, 1 ♀, Dingyuanying [Bayan Hot], Alashan [Helan Shan] Mt., 22–24.VI.1908, PK (ZISP); *Gansu*, 1 ♀, oasis Sachjou [Dunhuang] [40°09'N, 94°40'E], Gashun Gobi desert, 4.VIII.1895, leg. VR, PK (ZISP).

Distribution. *China (Inner Mongolia, Gansu), Central Asia, Mongolia, Russia, South Europe, Caucasus, Turkey, Iran, Israel, United Arab Emirates, North Africa, Cape Verde Islands.

Sphecodes scabricollis Wesmael, 1835

Figs 23, 40

Material examined. CHINA: *Heilongjiang*, 2 ♀♀, Harbin [45°46'N, 126°39'E], 11.VII.1954 (IZCAS); 1 ♀, idem, 8.VII.1955 (IZCAS); 1 ♀, idem, 19.VII.1955 (IZCAS); 1 ♀, idem, 20.VII.1955 (IZCAS); 2 ♀♀, idem, 25.VII.1955 (IZCAS); 1 ♂, idem, 8.VII.1955 (IZCAS), 2 ♂♂, idem, 25.VII.1955 (IZCAS); 1 ♂, idem, 11.IX.1955 (IZCAS); *Liaoning*, 1 ♂, Guicheng [43°40'N, 126°15'E], 10.VII.1962, leg. T.-L. Cheng (IZCAS); *Beijing*, 1 ♂, Changping district, Liucun town, Wangjiayuan village [40°12'N, 116°00'E], 5.IX.2011, leg. F. Yuan (IZCAS); 16 ♂♂, Wofosi [40°03'N, 115°10'E], 18.IX.1981, leg. Y.-R. Wu (IZCAS); 2 ♂♂, idem, 10.IX.1981, leg. Q. Zhou (IZCAS); 2 ♀♀, idem, 10.IX.1981, leg. W.-Z. Ma (IZCAS); 1 ♀, Bada Ling [40°22'N, 115°58'E], 3.IX.1981, leg. P.-Y. Yu (IZCAS); 1 ♂, idem, 28.VIII.1974 (IZCAS); 1 ♂, 13.VIII.1981 (IZCAS); 1 ♂, 20.VIII.1988, leg. Y.-S. Shi (IZCAS); 1 ♂, idem, 30.VIII.1977, leg. S.-F. Wang (IZCAS); 1 ♂, idem, 25.VIII.1981, leg. Q. Zhou (IZCAS); 1 ♀, idem, 7.IX.1982, leg. Z.-C. Jin (IZCAS); 1 ♂, Xiangshan



Figures 53–58. Antennae, males. **53** *Sphecoides pieli* Cockerell **54** *S. kozlovi* Astafurova & Proshchalykin **55** *S. pinguius* Pérez **56** *S. intermedius* Blüthgen **57** *S. gibbus* (Linnaeus) **58** *S. nippon* Meyer. Scale bars: 0.5 mm.

[39°54'N, 116°12'E], 22.IX.1983, leg. J.-G. Fan (IZCAS); 4 ♀♀, Qinglongqiao [39°54'N, 116°21'E], 5.IX.1988, leg. H.-L. Xu (IZCAS); 1 ♂, idem, 12.V.1981, leg. Y.-R. Wu (IZCAS); 4 ♀♀, 1 ♂, Beijing [39°55'N, 116°24'E], 28.VIII.1973, leg. Y.-R. Wu (IZCAS); 3 ♂♂, idem, 28.VIII.1973, leg. S.-F. Wang (IZCAS); *Shaanxi*, 2 ♀♀, Qihling Mts., 6 km E Xunyangba [33°32'N, 108°33'E], 1000–1300 m, 23.V–13.VI.1998, leg. Marshal (PCMS); *Qinghai*, 1 ♂, Sinin-khe River valley [36°30'N, 101°40'E], 29.VII.1908, leg. PK (ZISP); *Zhejiang*, 1 ♂, Lian Country, West Tianmu Mt. [30°20'N, 119°25'E], 1000 m, 16.IX.2000, S. Belokobylskij (ZISP).

Distribution. *China (Heilongjiang, Liaoning, Beijing, Shaanxi, Qinghai, Zhejiang), Kazakhstan (East Kazakhstan), Russia, Europe (north to S England and Latvia), Caucasus, Turkey, Iran, South Korea, Japan, India.

Sphecoides turanicus Astafurova & Proshchalykin, 2017

Figure 31

Material examined. CHINA: *Gansu*, 1 ♀, 2 ♂♂, oasis Sachjou [Dunhuang] [40°09'N, 94°40'E], Gashun Gobi desert, 1–9.VIII.1895, leg. VR, PK (ZISP).

Distribution. *China (Gansu), Central Asia.

Table 1. Checklist of the *Sphecodes* species of China including distribution by provinces.

	Species	Province	Published data	Type of areal
1	<i>S. albilabris</i> (Fabricius, 1793)	Gansu, Liaoning, Inner Mongolia, Shanxi	first record	P
2	<i>S. alternatus</i> Smith, 1853	Xinjiang, Gansu	first record	P
3	<i>S. chinensis</i> Meyer, 1922	China (exactly locality is unknown)	Meyer 1922	?
4	<i>S. crassus</i> Thomson, 1870	Inner Mongolia, Shanxi	first record	P
5	<i>S. cristatus</i> Hagens, 1882	Xinjiang, Inner Mongolia, Ningxia, Liaoning, Hebei, Shandong, Shanxi, Shanxi, Heilongjiang, Jilin, Beijing, Tianjin	Meyer 1922, Blüthgen 1927, Ascher and Pickering 2018, current data	P
6	<i>S. ephippius</i> (Linné, 1767)	Xinjiang	first record	P
7	<i>S. ferruginatus</i> Hagens, 1882	Shanxi, Beijing	first record	P
8	<i>S. formosanus</i> Cockerell, 1911	Taiwan	Cockerell 1911	O
9	<i>S. galeritus</i> Meyer, 1927	Guandong	Meyer 1927	O
10	<i>S. gibbus</i> (Linnaeus, 1758)	Xinjiang	Meyer 1920, current data	P
11	<i>S. geoffrellus</i> (Kirby, 1802)	Shanxi, Inner Mongolia	first record	P
12	<i>S. grahami</i> Cockerell, 1922	Sichuan, Shanghai; Hebei, Shaanxi, Shanxi, Jilin, Jiangsu, Anhui, Zhejiang, Xizang, Guandong, Yunnan	Cockerell 1922, 1931, Ascher and Pickering 2018, current data	PO
13	<i>S. howardi</i> Cockerell, 1922	Guandong	Cockerell 1922, Blüthgen 1924	O
14	<i>S. intermedius</i> Blüthgen, 1923	Gansu	first record	P
15	<i>S. kershawi</i> Perkins, 1921	Guandong	Meyer 1927	O
16	<i>S. kozlovi</i> Astafurova & Proshchalykin, 2015	Inner Mongolia, Ningxia, Shanxi	first record	P
17	<i>S. laticaudatus</i> Tsuneki, 1983	Hebei	first record	P
18	<i>S. laticeps</i> Meyer, 1920	Taiwan	Meyer 1920	O
19	<i>S. longulus</i> Hagens, 1882	Gansu, Shanxi, Hebei, Inner Mongolia	Blüthgen 1934, current data	P
20	<i>S. manchurianus</i> Strand & Yasumatsu, 1938	Liaoning	Strand and Yasumatsu 1938	P
21	<i>S. monilicornis</i> (Kirby, 1802)	Heilongjiang	first record	P
22	<i>S. nippon</i> Meyer, 1922	Gansu, Inner Mongolia, Shaanxi, Heilongjiang, Beijing, Gansu	Blüthgen 1934, current data	P
23	<i>S. nurekensis</i> Warncke, 1992	Xinjiang	first record	P
24	<i>S. olivieri</i> Lepeletier de Saint-Fargeau, 1825	Xinjiang, Gansu	first record	P
25	<i>S. pieli</i> Cockerell, 1931	Sichuan, Shanghai, Shanxi, Hebei, Beijing, Zhejiang, Jiangsu	Cockerell 1931, Ascher and Pickering 2018, current data	PO
26	<i>S. pinguiculus</i> Pérez, 1903	Gansu, Inner Mongolia	first record	P
27	<i>S. pectoralis</i> Morawitz, 1876	Xinjiang, Gansu	first record	P
28	<i>S. pellucidus</i> Smith, 1845	Xinjiang, Sichuan, Gansu	Blüthgen 1924, Meyer 1922, 1925, current data	P
29	<i>S. sauteri</i> Meyer, 1925	Taiwan	Meyer 1925	O
30	<i>S. scabricollis</i> Wesmael, 1835	Qinghai, Zhejiang, Shaanxi, Heilongjiang, Beijing	first record	P
31	<i>S. takaensis</i> Blüthgen, 1927	Taiwan	Blüthgen 1927	O
32	<i>S. tertius</i> Blüthgen, 1927	Guandong	Ascher and Pickering 2018	O
33	<i>S. turanicus</i> Astafurova & Proshchalykin, 2017	Gansu	first record	P

P – Palearctic species; O – Oriental species; PO – Palearctic and Oriental species

Discussion

In total, 33 species of *Sphecodes* are recorded from China (Table 1). Twenty-two of these species are Palearctic and two species have a Palearctic-Oriental range. For comparison, 37 species are known from Russia, 20 species of these from the Russian Far East (Astafurova and Proshchalykin 2014, 2017b) and 36 from Central Asia (Astafurova and Proshchalykin 2017a, Astafurova et al. 2018). In contrast, the Oriental fauna of the genus is poorly studied: only nine species are recorded from Oriental China, most of which are only known from type series, suggesting that further revision is necessary.

The majority of the Palearctic Chinese *Sphecodes* is composed of 14 widespread Trans-Palearctic or Euro-Asian species. Of them, eight species are distributed from Europe to the Russian Far East, Japan and the eastern provinces of China (*S. albilabris*, *S. crassus*, *S. cristatus*, *S. ferruginatus*, *S. geoffrellus*, *S. longulus*, *S. monilicornis*, and *S. scabricollis*). One species, *S. pellucidus*, occurs in the Russian Far East, but is rare in the East Palearctic and has not yet been found in eastern China. Five species are distributed from Europe to Siberia and are, as expected, recorded in north-west China (*S. alternatus*, *S. gibbus*, *S. ephippius*, *S. pinguiculus*, *S. intermedius*).

The other eight Palearctic species have smaller distributional ranges. *Sphecodes olivieri* is found in semi-desert and desert habitats of the Western Palearctic, including Xinjiang and Gansu within China. Three species, *S. nurekensis*, *S. pectoralis* and *S. turanicus*, are desert and steppe Irano-Turanian species distributed in Central Asia which also occur in Xinjiang and Gansu. Four species, *S. kozlovi*, *S. laticaudatus*, *S. manchurianus*, and *S. nippon* are East Palearctic species, not found farther west than 90°E.

None of the above Palearctic species are recorded below 30°N. However, two species, *S. grahami* and *S. pieli*, have an inter-realm range and are distributed in the East Palearctic well as Oriental China. More study is necessary to revise the Oriental *Sphecodes*.

Acknowledgments

We are grateful to Maximilian Schwarz and Fritz Gusenleitner (OÖLM) for help during our visit to Austria. We thank Hege Vårdal (NHRS), Michael Ohl and Viola Richter (MNHBB) for providing *Sphecodes* type specimens. Michael Orr helped to edit the English text during his busy field trips. We also wish to thank Peter Bogusch and Julio Genaro, who reviewed the manuscript and helped to improve this article. This investigation was supported by the Russian Funds for Basic Research (grant numbers 16–04–00197 and 17–04–00259) and the state research project (AAAA-A17–117030310210–3). Ze-Qin Niu is supported by the National Science Foundation of China (31772487). Chao-Dong ZHU acknowledges the supports of the National Science Fund for Distinguished Young Scholars (Grant number 31625024) and Singapore-China Joint Research Grant (41761144068) to work on bee systematics.

References

- Årgent L, Svensson B (1982) Flagellar sensilla on *Sphecodes* bees (Hymenoptera, Halictidae). *Zoologica Scripta* 11: 45–54. <https://doi.org/10.1111/j.1463-6409.1982.tb00517.x>
- Ascher JS, Pickering J (2018) Discover Life bee species guide and world checklist (Hymenoptera: Apoidea: Anthophila). http://www.discoverlife.org/mp/20q?guide=Apoidea_species [Accessed 25 May 2018]
- Astafurova YuV (2013) Geographic distribution of halictid bees of the subfamilies Rophitinae and Nomiinae (Hymenoptera, Halictidae) in the Palaearctic. *Entomological Review* 93(4): 437–451. <https://doi.org/10.1134/S0013873813040052>
- Astafurova YuV, Proshchalykin MYu (2014) The bees of the genus *Sphecodes* Latreille 1804 of the Russian Far East, with key to species (Hymenoptera: Apoidea: Halictidae). *Zootaxa* 3887(5): 501–528. <http://dx.doi.org/10.11646/zootaxa.3887.5.1>
- Astafurova YuV, Proshchalykin MYu (2015a) Bees of the genus *Sphecodes* Latreille 1804 of Siberia, with a key to species (Hymenoptera: Apoidea: Halictidae). *Zootaxa* 4052(1): 65–95. <http://dx.doi.org/10.11646/zootaxa.4052.1.3>
- Astafurova YuV, Proshchalykin MYu (2015b) New and little known bees of the genus *Sphecodes* Latreille (Hymenoptera: Halictidae) from Mongolia. *Far Eastern Entomologist* 289: 1–9.
- Astafurova YuV, Proshchalykin MYu (2016a) To the knowledge of the genus *Sphecodes* Latreille (Hymenoptera: Halictidae) of Caucasus. *Euroasian Entomological Journal* 5(Suppl. 1): 15–9.
- Astafurova YuV, Proshchalykin MYu (2016b) The bees of the genus *Sphecodes* Latreille (Hymenoptera: Halictidae) of the European part of Russia. *Far Eastern Entomologist* 321: 1–21.
- Astafurova YuV, Proshchalykin MYu (2017a) To the knowledge of the *Sphecodes hyalinatus* species-group (Hymenoptera, Halictidae). *Entomological Review* 97(5): 664–671. <https://doi.org/10.1134/S0013873817050104>
- Astafurova YuV, Proshchalykin MYu (2017b) The genus *Sphecodes* Latreille 1804 (Hymenoptera: Apoidea: Halictidae) in Central Asia. *Zootaxa* 4324(2): 249–284. <https://doi.org/10.11646/zootaxa.4324.2.3>
- Astafurova YuV, Proshchalykin MYu, Schwarz M (2015) New data on the genus *Sphecodes* Latreille (Hymenoptera: Halictidae) from Mongolia. *Far Eastern Entomologist* 302: 1–9.
- Astafurova YuV, Proshchalykin MYu, Schwarz M (2018) New and little known bees of the genus *Sphecodes* Latreille, 1804 (Hymenoptera: Apoidea: Halictidae) from Central Asia. *Zootaxa* 4441(1): 76–88. <https://doi.org/10.11646/zootaxa.4441.1.4>
- Astafurova YuV, Proshchalykin MYu, Shlyakhtenok AS (2014) Contribution to the knowledge of bee fauna of the genus *Sphecodes* Latreille (Hymenoptera: Halictidae) of the Republic of Belarus. *Far Eastern Entomologist* 280: 1–8.
- Blüthgen P (1923) Beiträge zur Systematik der Bienengattung *Sphecodes* Latr. *Deutsche Entomologische Zeitschrift* 5: 441–513.
- Blüthgen P (1924) Beiträge zur Systematik der Bienengattung *Sphecodes* Latr. II. *Deutsche Entomologische Zeitschrift* 6: 457–516.
- Blüthgen P (1927) Beiträge zur Systematik der Bienengattung *Sphecodes* Latr. III. *Zoologische Jahrbücher. Abteilung für Systematik, Ökologie und Geographie der Tiere* 53(1/3): 23–112.

- Blüthgen P (1934) Schwedisch-chinesische wissenschaftliche Expedition nach den nordwestlichen Provinzen Chinas unter Leitung von Dr. Sven Hedin und Prof. Sü Ping-chang: Insekten, gesammelt vom schwedischen Arzt der Expedition Dr. David Hummel 1927–1930. 27 Hymenoptera. 5. *Halictus*- und *Sphecodes*-Arten (Hym.; Apidae; Halictini). Arkiv för Zoologi 27(1, 13): 1–23.
- Bogusch P, Straka J (2012) Review and identification of the cuckoo bees of central Europe (Hymenoptera: Halictidae: *Sphecodes*). Zootaxa 3311: 1–41.
- Chen L, Song Y, Xu S (2008) The boundary of Palaearctic and Oriental realms in western China. Progress in Natural Science 18: 833–841. <https://doi.org/10.1016/j.pnsc.2008.02.004>
- Cockerell TDA (1911) Descriptions and records of bees – XXXIV. Annals and Magazine of Natural History 7(39): 225–237. <https://doi.org/10.1080/00222931108692933>
- Cockerell TDA (1922) Bees in the collection of the United States National Museum. 4. Proceedings of the United States National Museum 60(2413): 1–20. <https://doi.org/10.5479/si.00963801.60-2413.1>
- Cockerell TDA (1931) Bees collected by the Reverend O. Piel in China. American Museum Novitates 466: 1–16.
- Emeljanov AF (1974) Proposals on Classification and Nomenclature of Ranges. Entomologicheskoe Obozrenie 53(3): 497–521. [In Russian]
- Engel MS (2001) A monograph of the Baltic amber bees and evolution of the Apoidea (Hymenoptera). Bulletin of the American Museum of Natural History 259: 1–192. [https://doi.org/10.1206/0003-0090\(2001\)259<0001:AMOTBA>2.0.CO;2](https://doi.org/10.1206/0003-0090(2001)259<0001:AMOTBA>2.0.CO;2)
- Fellowes JR (2006) Ant (Hymenoptera: Formicidae) genera in southern China: observations on the Oriental-Palaearctic boundary. Myrmecologische Nachrichten 8: 239–249.
- He J, Kreft H, Gao E, Wang Z, Jiang H (2017) Patterns and drivers of zoogeographical regions of terrestrial vertebrates in China. Journal of Biogeography 44: 1172–1184. <https://doi.org/10.1111/jbi.12892>
- Heiser M, Schmitt T (2013) Tracking the boundary between the Palaearctic and the Oriental region: new insights from dragonflies and damselflies (Odonata). Journal of Biogeography 40: 2047–2058. <https://doi.org/10.1111/jbi.12133>
- Hoffmann RS (2001) The Southern boundary of the Palaearctic realm in China and adjacent countries. Acta Zoologica Sinica 47(2): 121–131.
- Meyer R (1920) Apidae Sphecodinae. Archiv für Naturgeschichte. Abteilung A 85(1/2): 79–160; 161–244.
- Meyer R (1922) Nachtrag 1 zur Bienengattung *Sphecodes* Latr. Archiv für Naturgeschichte. Abteilung A 88(8): 165–174.
- Meyer R (1925) Zur Bienengattung *Sphecodes*. Archiv für Naturgeschichte. Abteilung A 90(12): 1–12.
- Michener CD (2007) The Bees of the World (2nd edn). Johns Hopkins University Press, Baltimore, 953 pp.
- Pesenko YuA (2007) Family Halictidae. In: Lelej AS (Ed.) Key to the insects of Russian Far East. Volume 4. Part 5. Dalnauka, Vladivostok, 745–878. [In Russian]
- Strand E, Yasumatsu K (1938) Two new species of the genus *Sphecodes* Latreille from the Far East (Hymenoptera: Apoidea). Mushi 11(1): 78–82.
- Warncke K (1992) Die westpaläarktischen Arten der Bienengattung *Sphecodes* Latr. Bericht der Naturforschende Gesellschaft Augsburg 52: 9–64.